

Relevante Studien zur Behandlung von Verbaler Entwicklungsdypraxie, die **keine** RCTs sind

übernommen aus dem Paper: Morgan, A. T., Murray, E., & Liegeois, F. J. (2018). Interventions for childhood apraxia of speech. *Cochrane Database of Systematic Reviews*, (5).

VED-Therapiestudien

- Baas BS, Strand EA, LM Elmer, Barbaresi WJ. Treatment of severe childhood apraxia of speech in a 12-year-old male with CHARGE association. *Journal of Medical Speech-Language Pathology* 2008;16(4):181-90.
- Ballard KJ, Robin DA, McCabe P, McDonald J. A treatment for dysprosody in childhood apraxia of speech. *Journal of Speech, Language, and Hearing Research* 2010;53(5):1227-45. [DOI: 10.1044/1092-4388(2010/09-0130); PUBMED: 20798323]
- Beathard B, Krout RE. A music therapy clinical case study of a girl with childhood apraxia of speech: finding Lily's voice. *Arts in Psychotherapy* 2008;35(2):107-16. [DOI: 10.1016/j.aip.2008.01.004]
- Binger C, Light J. The effect of aided AAC modeling on the expression of multi-symbol messages by preschoolers who use AAC. *Augmentative and Alternative Communication* 2007;23(1):30-43. [DOI: 10.1080/07434610600807470; PUBMED: 17364486]
- Binger C, Light J. The morphology and syntax of individuals who use AAC: research review and implications for elective practice. *Augmentative and Alternative Communications* 2008;24(2):123-38. [DOI: 10.1080/07434610701830587; PUBMED: 18465366]
- Binger C, Maguire-Marshall M, Kent-Walsh J. Using aided AAC models, recasts, and contrastive targets to teach grammatical morphemes to children who use AAC. *Journal of Speech, Language, and Hearing Research* 2011;54(1):160-76. [DOI: 10.1044/1092-4388(2010/09-0163); PUBMED: 20719874]
- Bornman J, Alant E, Meiring E. The use of a digital voice output device to facilitate language development in a child with developmental apraxia of speech: a case study. *Disability and Rehabilitation* 2001;23(14):623-34. [PUBMED: 11697460]
- Bose A, Square PA, Schlosser R, van Lieshout P. Effects of PROMPT therapy on speech motor function in a person with aphasia and apraxia of speech. *Aphasiology* 2001;15(8):767-85. [DOI: 10.1080/02687040143000186]
- Carter P, Edwards S. EPG therapy for children with long-standing speech disorders: predictions and outcomes. *Clinical Linguistics and Phonetics* 2004;18(6-8):359-72. [PUBMED: 15573477]
- Chappell GE. Childhood verbal apraxia and its treatment. *Journal of Speech and Hearing Disorders* 1973;38(3):362-8. [PUBMED: 4721796]
- Culp DM. Developmental apraxia and augmentative or alternative communication - a case example. *Augmentative and Alternative Communication* 1989;5(1):27-34. [DOI: 10.1080/07434618912331274936]
- Cumley GD, Swanson S. Augmentative and alternative communication options for children with developmental apraxia of speech: three case studies. *Augmentative and Alternative Communication* 1999;15(2):110-25. [DOI: 10.1080/07434619912331278615]

Dale PS, Hayden DA. Treating speech subsystems in childhood apraxia of speech with tactual input: the PROMPT approach. *American Journal of Speech-language Pathology* 2013;22(4):644-61. [DOI: 10.1044/1058-0360(2013/12-0055); PUBMED: 23813194]

Daly DA, Cantrell RP, Cantrell ML, Aman LA. Structure speech therapy contingencies with an oral apraxic child. *Journal of Speech and Hearing Disorders* 1972;37(1):22-32. [DOI: 10.1044/jshd.3701.22]

Dworkin JP, Abkarian GG, John DF. Apraxia of speech: the effectiveness of a treatment regimen. *Journal of Speech and Hearing Disorders* 1988;53(3):280-94. [PUBMED: 3398481]

Edeal DM, Gildersleeve-Neumann CE. The importance of production frequency in therapy for childhood apraxia of speech. *American Journal of Speech-language Pathology* 2011;20(2):95-110. [DOI: 10.1044/1058-0360(2011/09-0005); PUBMED: 21330650]

Forrest K, Elbert M. Treatment for phonologically disordered children with variable substitution patterns. *Clinical Linguistics & Phonetics* 2001;15(1-2):41-5. [DOI:10.3109/02699200109167628; PUBMED: 21269096]

Groenen P, Maassen B, Crul T, Thoonen G. The specific relation between perception and production errors for place of articulation in developmental apraxia of speech. *Journal of Speech and Hearing Research* 1996;39(3):468-82. [PUBMED: 8783127]

Hadar U, Twiston-Davies R, Steiner TJ, Rose FC. A psychomotor approach to improving speech by modulating suprasegmental control in motor dysphasia and articulatory apraxia. *Advances in Neurology* 1984;42:337-51. [PUBMED: 6507181]

Hall PK. The occurrence of developmental apraxia of speech in a mild articulation disorder: a case study. *Journal of Communication Disorders* 1989;22(4):265-76. [PUBMED: 2794108]

Hall PK, Hardy JC, LaVelle WE. A child with signs of developmental apraxia of speech with whom a palatal liM prosthesis was used to manage palatal dysfunction. *Journal of Speech and Hearing Disorders* 1990;55(3):454-60. [PUBMED: 2381187]

Harris L, Doyle ES, Haaf R. Language treatment approach for users of AAC: experimental single-subject investigation. *Augmentative and Alternative Communication* (Baltimore, Md. : 1985) 1996;12(4):230-43. [DOI: 10.1080/07434619612331277698]

Hayden D. The PROMPT model: use and application for children with mixed phonological-motor impairment. *Advances in Speech Language Pathology* 2006;8(3):265-81. [DOI: 10.1080/14417040600861094]

Hayden D. The PROMPT model: use and application for children with mixed phonological-motor impairment. *Advances in Speech Language Pathology* 2006;8(3):265-81. [DOI: 10.1080/14417040600861094]

Head DG, Smith D. Speech remediation of children involved in two different physical education programs. *Perceptual and Motor Skills* 1975;40(1):261-2. [DOI: 10.2466/pms.1975.40.1.261; PUBMED: 1118271]

Helfrich-Miller KR. A clinical perspective: melodic intonation therapy for developmental apraxia. *Clinics in Communication Disorders* 1994;4(3):175-82. [PUBMED: 7994292]

Iuzzini J, Forrest K. Evaluation of a combined treatment approach for childhood apraxia of speech. *Clinical Linguistics and Phonetics* 2010;24(4-5):335-45. [DOI: 10.3109/02699200903581083; PUBMED: 20345262]

Jaroma M, Danner P, Koivuniemi E. Sensory integrative therapy and speech therapy for improving the perceptual motor skills and speech articulation of a dyspractic boy. *Folia Phoniatrica* 1984;36(6):261-6. [DOI: 10.1159/000265753]

Kadis DS, Goshulak D, Namasivayam A, Pukonen M, Kroll R, De Nil LF, et al. Cortical thickness in children receiving intensive therapy for idiopathic apraxia of speech. *Brain Topography* 2014;27(2):240-7. [DOI: 10.1007/s10548-013-0308-8; PMC3921462; PUBMED: 23974724]

Katz WF, Bharadwaj SV, Stettler MP. Influences of electromagnetic articulography sensors on speech produced by healthy adults and individuals with aphasia and apraxia. *Journal of Speech, Language, and Hearing Research* 2006;49(3):645-59. [DOI: 10.1044/1092-4388(2006/047); PUBMED: 16787902]

King AM, Hengst JA, DeThorne LS. Severe speech sound disorders: an integrated multimodal intervention. *Language, Speech, and Hearing Services in Schools* 2013;44(2):195-210. [DOI: 10.1044/0161-1461(2012/12-0023); PUBMED: 23633644]

Kingston LM, Rosenthal JB. Oral stereognosis in children with disordered articulation: measurement issues, and a treatment study. *Australian Journal of Human Communication Disorders* 1987;15(1):1-14. [DOI: 10.3109/asl2.1987.15.issue-1.01]

Klick SL. Adapted cuing technique for use in treatment of dyspraxia. *Language, Speech, and Hearing Services in Schools* 1985;16:256-9. [DOI: 10.1044/0161-1461.1604.256]

Krauss T, Galloway H. Melodic intonation therapy with language delayed apraxic children. *Journal of Music Therapy* 1982;19(2):102-13. [DOI: 10.1093/jmt/19.2.102]

Lagasse B. Evaluation of melodic intonation therapy for developmental apraxia of speech. *Music Therapy Perspectives* 2012;30(1):49-55. [DOI: 10.1093/mtp/30.1.49]

Lozano RA, Dreyer DE. Some effects of delayed auditory feedback on dyspraxia of speech. *Journal of Communication Disorders* 1978;11(5):407-15. [PUBMED: 730833]

Lüke C. Impact of speech-generating devices on the language development of a child with childhood apraxia of speech: a case study. *Disability and Rehabilitation. Assistive Technology* 2016;11(1):80-8. [DOI: 10.3109/17483107.2014.913715; PUBMED: 24773213]

Lundeborg I, McAllister A. Treatment with a combination of intra-oral sensory stimulation and electropalatography in a child with severe developmental dyspraxia. *Logopedics, Phoniatrics, Vocology* 2007;32(2):71-9. [DOI: 10.1080/14015430600852035; PUBMED: 17613788]

Maas E, Butalla CE, Farinella KA. Feedback frequency in treatment for childhood apraxia of speech. *American Journal of Speech-language Pathology* 2012;21(3):239-57. [DOI:10.1044/1058-0360(2012/11-0119); PUBMED: 22442284]

Martikainen 2011 {published data only} Martikainen A-L, Korpilahti P. Intervention for childhood apraxia of speech: a single-case study. *Child Language Teaching and Therapy* 2011;27(1):9-20. [DOI: 10.1177/0265659010369985]

Martin MK, Wright LE, Perry S, Cornett D, Schraeder M, Johnson JT. Children with developmental verbal dyspraxia: changes in articulation and perceived resilience with intensive multimodal intervention. *Child Language Teaching and Therapy* 2016;32(3):261-75. [DOI: 10.1177/0265659015615924]

McCabe P, Macdonald-D'Silva AG, van Rees LJ, Ballard KJ, Arciuli J. Orthographically sensitive treatment for dysprosody in children with childhood apraxia of speech using ReST intervention. *Developmental Neurorehabilitation* 2014;17(2):137-46. [DOI: 10.3109/17518423.2014.906002; PUBMED: 24694312]

McNeill BC, Gillon GT, Dodd B. Phonological awareness and early reading development in childhood apraxia of speech (CAS). *International Journal of Language & Communication Disorders / Royal College of Speech & Language Therapists* 2009;44(2):175-92. [DOI: 10.1080/13682820801997353; PUBMED: 19234970]

McNeill BC, Gillon GT, Dodd B. A longitudinal case study of the effects of an integrated phonological awareness program for identical twin boys with childhood apraxia of speech (CAS). *International Journal of Speech-language Pathology* 2009;11(6):482-95. [DOI: 10.3109/17549500902842583; PUBMED: 21271925]

- McNeill BC, Gillon GT, Dodd B. The longer term effects of an integrated phonological awareness intervention for children with childhood apraxia of speech. *Asia Pacific Journal of Speech, Language, and Hearing* 2010;13(3):145-61. [DOI: 10.1179/136132810805335074]
- Morgan Barry R. The relationship between dysarthria and verbal dyspraxia in children: a comparative study using profiling and instrumental analyses. *Clinical Linguistics & Phonetics* 1995;9(4):277-309. [DOI: 10.3109/02699209508985338]
- Moriarty BC, Gillon GT. Phonological awareness intervention for children with childhood apraxia of speech. *International Journal of Language & Communication Disorders / Royal College of Speech & Language Therapists* 2006;41(6):713-34. [DOI: 10.1080/13682820600623960; PUBMED: 17079224]
- Namasivayam AK, Pukonen M, Goshulak D, Yu VY, Kadis DS, Kroll R, et al. Relationship between speech motor control and speech intelligibility in children with speech sound disorders. *Journal of Communication Disorders* 2013;46(3):264-80. [DOI: 10.1016/j.jcomdis.2013.02.003; PUBMED: 23628222]
- Namasivayam A, Pukonen M, Hard J, Jahnke R, Kearney E, Kroll R, et al. Motor speech treatment protocol for developmental motor speech disorders. *Developmental Neurorehabilitation* 2015;18(5):296-303. [DOI: 10.3109/17518423.2013.832431; PUBMED: 24088085]
- Preston JL, Brick N, Landi N. Ultrasound biofeedback treatment for persisting childhood apraxia of speech. *American Journal of Speech-language Pathology* 2013;22(4):627-43. [DOI: 10.1044/1058-0360(2013/12-0139); PUBMED: 23813207]
- Preston JL, Leece MC, Maas E. Intensive treatment with ultrasound visual feedback for speech sound errors in childhood apraxia. *Frontiers in Human Neuroscience* 2016;10:440. [DOI: 10.3389/fnhum.2016.00440; PMC5003919; PUBMED: 27625603]
- Preston JL, Leece MC, Maas E. Motor-based treatment with and without ultrasound feedback for residual speech-sound errors. *International Journal of Language and Communication Disorders* 2017;52(1):80-94. [DOI: 10.1111/1460-6984.12259; PMC5156595; PUBMED: 27296780]
- Ray J. Effects of orofacial myofunctional therapy on speech intelligibility in individuals with persistent articulatory impairments. *International Journal of Orofacial Myology* 2003;29:5-14. [PUBMED: 14689652]
- Richardson AJ. Clinical trials of fatty acid treatment in ADHD, dyslexia, dyspraxia and the autistic spectrum. *Prostaglandins, Leukotrienes, and Essential Fatty Acids* 2004;70(4):383-90. [DOI: 10.1016/j.plefa.2003.12.020; PUBMED: 15041031]
- Rosenbek J, Hansen R, Baughman CH, Lemme M. Treatment of developmental apraxia of speech: a case study. *Language, Speech, and Hearing Services in Schools* 1974;5:13-22. [DOI: 10.1044/0161-1461.0501.13]
- Rosenthal JB. Rate control therapy for developmental apraxia of speech. *Clinics in Communication Disorders* 1994;4(3):190-200. [PUBMED: 7994294]
- Skelton SL, Hagopian AL. Using randomized variable practice in the treatment of childhood apraxia of speech. *American Journal of Speech-language Pathology* 2014;23(4):599-611. [DOI: 10.1044/2014_AJSLP-12-0169; MEDLINE: 25017177]
- Square PA. Treatment approaches for developmental apraxia of speech. *Clinics in Communication Disorders* 1994;4(3):151-61. [PUBMED: 7994290]
- Stokes SF, GriJiths R. The use of facilitative vowel contexts in the treatment of post-alveolar fronting: a case study. *International Journal of Language & Communication Disorders / Royal College of Speech & Language Therapists* 2010;45(3):368-80. [DOI: 10.3109/13682820903094737; PUBMED: 20144008]
- Strand EA, Debertine P. The efficacy of integral stimulation intervention with developmental apraxia of speech. *Journal of Medical Speech-language Pathology* 2000;8(4):295-300. [www.researchgate.net/publication/286964810_The_efficacy_of_integral_stimulation_intervention_with_developmental_apraxia_of_speech]

- Strand EA, Stoeckel R, Baas B. Treatment of severe childhood apraxia of speech: a treatment efficacy study. *Journal of Medical Speech-language Pathology* 2006;14(4):297-307. [psycnet.apa.org/record/2006-22884-013]
- Thomas DC, McCabe P, Ballard KJ. Rapid Syllable Transitions (ReST) treatment for childhood apraxia of speech: the effect of lower dose-frequency. *Journal of Communication Disorders* 2014;51:29-42. [DOI: 10.1016/j.jcomdis.2014.06.004; PUBMED: 25052390]
- Thomas DC, McCabe P, Ballard KJ, Lincoln M. Telehealth delivery of Rapid Syllable Transitions (ReST) treatment for childhood apraxia of speech. *International Journal of Language & Communication Disorders* 2016;51(6):654-71. [DOI: 10.1111/1460-6984.12238; PUBMED: 27161038]
- Tierney CD, Pitterle K, Kurtz M, Nakhla M, Todorow C. Bridging the gap between speech and language: using multimodal treatment in a child with apraxia. *Pediatrics* 2016;138(3):e20160007. [DOI: 10.1542/peds.2016-0007; PUBMED: 27492818]
- Vashdi E. Using VML (verbal motor learning) method techniques in treatment of prosody disorder due to childhood apraxia of speech: a case study. *International Journal of Child Health and Human Development* 2013;6(2):255-60. [www.novapublishers.com/catalog/product_info.php?products_id=52658]
- Vashdi E. The influence of initial phoneme cue technique on word formation: a case study of a child with apraxia of speech and autism. *International Journal of Child Health and Human Development* 2014;7(2):197-203. [www.novapublishers.com/catalog/product_info.php?products_id=53974]
- Velleman SL, Strand K. Developmental verbal dyspraxia. In: Bernthal JE, Bankson NW editor(s). *Child Phonology: Characteristics, Assessment, and Intervention with Special Populations*. New York (NY): Thieme, 1994:110-39. [ISBN0-86577-502-8]
- Yoss KA, Darley FL. Developmental apraxia of speech in children with defective articulation. *Journal of Speech and Hearing Research* 1974;17(3):399-416. [PUBMED: 4421901]
- Zaretsky E, Velleman SL, Curro K. Through the magnifying glass: underlying literacy deficits and remediation potential in childhood apraxia of speech. *International Journal of Speech-language Pathology* 2010;12(1):58-68. [PUBMED: 20380250]

Weitere Quellen für VED-Therapie und verwandte Themen

- American Speech-Language-Hearing Association (ASHA). Technical Report. Childhood Apraxia of Speech: Ad Hoc Committee on Apraxia of Speech in Children. www.asha.org/policy/TR2007-00278/ (accessed 20 April 2018).
- Chumpelik D. The PROMPT system of therapy: theoretical framework and applications for developmental apraxia of speech. *Seminars in Speech and Language* 1984;5(2):139-56. [DOI: 10.1055/s-0028-1085172]
- Crosbie S, Holm A, Dodd B. Intervention for children with severe speech disorder: a comparison of two approaches. *International Journal of Language and Communication Disorders* 2005;40(4):469-71. [DOI: 10.1080/13682820500126049; PUBMED: 16195201]
- Davis BL, Jacks A, Marquardt TP. Vowel patterns in developmental apraxia of speech: three longitudinal casestudies. *Clinical Linguistics and Phonetics* 2005;19(4):249-74. [DOI: 10.1080/02699200410001695367; PUBMED: 16019775]
- Delaney AL, Kent RD. Developmental profiles of children diagnosed with apraxia of speech. Annual Convention of the American Speech-Language-Hearing Association; 2004 Nov 18-20; Philadelphia (PA). 2004.
- Dodd B, Hua Z, Crosbie S, Holm A, Ozanne A. DEAP: Diagnostic Evaluation of Articulation and Phonology. San Antonio (TX): PsychCorp of Harcourt Assessment, 2006.
- Dodd B, Crosbie S, McIntosh B, Holm A, Harvey C, Liddy M, et al. The impact of selecting di(er)ence contrasts in phonological therapy. *International Journal of Speech-Language Pathology* 2008;10(5):334-45. [DOI: 10.1080/14417040701732590; PUBMED: 20840033]
- Eadie P, Morgan A, Ukoumunne OC, Ttofari Eecen K, Wake M, Reilly S. Speech sound disorder at 4 years: prevalence, comorbidities, and predictors in a community cohort of children. *Developmental Medicine and Child Neurology* 2015;57(6):578-84. [DOI: 10.1111/dmcn.12635; PUBMED: 25403868]
- Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315(7109):629-34. [DOI: 10.1136/bmj.315.7109.629; PMC2127453; PUBMED: 9310563]
- Eising E, Carrion-Castillo A, Vino A, Strand EA, Jakielski KJ, Scerri TS, et al. A set of regulatory genes co-expressed in embryonic human brain is implicated in disrupted speech development. *Molecular Psychiatry* 2018 Feb 20 [Epub ahead of print]. [DOI: 10.1038/s41380-018-0020-x; PUBMED: 29463886]
- Fedorenko E, Morgan A, Murray E, Cardinaux A, Mei C, Tager-Flusberg H, et al. A highly penetrant form of childhood apraxia of speech due to deletion of 16p11.2. *European Journal of Human Genetics* 2016;24(2):302-6. [DOI: 10.1038/ejhg.2015.149; PMC4717199; PUBMED: 26173965]
- Gillon GT. The efficacy of phonological awareness intervention for children with spoken language impairment. *Language, Speech and Hearing Services in Schools* 2000;31(2):126-41. [DOI: 10.1044/0161-1461.3102.126; PUBMED: 27764385]
- Goldman R, Fristoe M. Goldman-Fristoe Test of Articulation-2. 2nd Edition. Minneapolis (MN): Pearson Assessments, 2000.
- Jacks A, Marquardt TP, Davis BL. Consonant and syllable structure patterns in childhood apraxia of speech: developmental change in three children. *Journal of Communication Disorders* 2006; Vol. 39, issue 6:424-41. [DOI: 10.1016/j.jcomdis.2005.12.005]
- Lai CS, Fisher SE, Hurst JA, Vargha-Khadem F, Monaco AP. A forkhead-domain gene is mutated in a severe speech and language disorder. *Nature* 2001;413(6855):519-23. [DOI: 10.1038/35097076; PUBMED: 11586359]
- Liégeois FJ, Morgan AT. Neural bases of childhood speech disorders: lateralization and plasticity for speech functions during development. *Neuroscience and Biobehavioral Reviews* 2012;36(1):439-58. [DOI: 10.1016/j.neubiorev.2011.07.011; PUBMED: 21827785]

- Liégeois F, Mayes A, Morgan A. Neural correlates of developmental speech and language disorders: evidence from neuroimaging. *Current Developmental Disorders Reports* 2014;1(3):215-27. [DOI: 10.1007/s40474-014-0019-1; PUBMED: PMC4104164]
- Maas E, Robin DA, Austermann Hula SN, Freedman SE, Wulf G, Ballard KJ, et al. Principles of motor learning in treatment of motor speech disorders. *American Journal of Speech-language Pathology* 2008;17(3):277-98. [DOI: 10.1044/1058-0360(2008/025); PUBMED: 18663111]
- Maas E, Gildersleeve-Neumann CE, Jakielski KJ, Stoeckel R. Motor-based intervention protocols in treatment of childhood apraxia of speech (CAS). *Current Developmental Disorders Reports* 2014;1(3):197-206. [DOI: 10.1007/s40474-014-0016-4; PMC4192721; PUBMED: 25313348]
- McNeill BC, Gillon GT, Dodd B. Effectiveness of an integrated phonological awareness approach for children with childhood apraxia of speech (CAS). *Child Language Teaching and Therapy* 2009;25(3):341-66. [DOI: 10.1177/0265659009339823]
- Morgan A, Fisher SE, ScheJér IE, Hildebrand M. FOXP2-related speech and language disorders. 2016 Jun 23 [updated 2017 Feb 2]. In: Adam MP, Ardinger HH, Pagon RA, Wallace SE, Bean LJH, Stephens K, et al. editor(s). *GeneReviews®*. Seattle (WA): University of Washington, 1993-2018. [www.ncbi.nlm.nih.gov/books/NBK368474/?report=reader#_NBK368474_pubdet]
- Morgan AT, van HaaMen L, van Hulst K, Edley C, Mei C, Yang Tan T, et al. Early speech development in Koolen de Vries Syndrome limited by oral praxis and hypotonia. *European Journal of Human Genetics* 2018; Vol. 26, issue 1:75-84. [DOI: 10.1038/s41431-017-0035-9; PUBMED: 29225339]
- Murray E, McCabe P, Ballard KJ. A comparison of two treatments for childhood apraxia of speech: methods and treatment protocol for a parallel group randomised control trial. *BMC Pediatrics* 2012;12:112. [DOI: 10.1186/1471-2431-12-112; ACTRN12612000744853; PMC3441276; PUBMED: 22863021]
- Royal College of Speech and Language Therapists (RCSLT). Developmental verbal dyspraxia policy statement. www.rcslt.org/speech_and_language_therapy/rcslt_position_papers (accessed prior to 21 March 2018). Review Manager 2014 [Computer program] Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager 5 (RevMan 5). Version 5.3. Copenhagen: Nordic Cochrane Centre, The Cochrane Collaboration, 2014.
- Shriberg LD, Potter NL, Strand EA. Prevalence and phenotype of childhood apraxia of speech in youth with galactosemia. *Journal of Speech, Language, and Hearing Research* 2011;54(2):487-519. [DOI: 10.1044/1092-4388(2010/10-0068); PMC3070858; PUBMED: 20966389]
- Turner SJ, Mayes AK, Verhoeven A, Mandelstam SA, Morgan AT, ScheJér IE. GRIN2A: an aptly named gene for speech dysfunction. *Neurology* 2015;84(6):586-93. [DOI: 10.1212/WNL.0000000000001228; PMC4335991; PUBMED: 25596506]
- Williams P, Stephens H, editor(s). *The Nuffield Centre Dyspraxia Programme*. 3rd Edition. London (UK): The Nuffield Centre Dyspraxia Programme Ltd., 2004. [www.ndp3.org]
- Williams P, Stephens H. *The Nuffield Centre Dyspraxia Programme*. In: Williams AL, McLeod S, McAuley RJ editor(s). *Interventions for Speech Sound Disorders in Children*. Baltimore (MD): Brookes Publishing Company, 2010:159-77.
- Yoss KA. Developmental apraxia of speech in children: familial patterns and behavioral characteristics. *American Speech and Hearing Association North Central Regional Conference*, 1975 May 9; Minneapolis (MN).

Die Literaturangaben sind übernommen aus dem Paper Morgan, A. T., Murray, E., & Liegeois, F. J. (2018). Interventions for childhood apraxia of speech. *Cochrane Database of Systematic Reviews*, (5). als Zusatzmaterial für #61 LingoScience, dem Logostudienpodcast.

Lingo Lab – Digitale Lösungen für die Sprachtherapie www.lingo-lab.de

dbS – Deutscher Bundesverband für akademische Sprachtherapie und Logopädie www.dbS-ev.de